



LETTER

PML-RARA mutations confer varying arsenic trioxide resistance

Dear Editor,

Acute promyelocytic leukemia (APL) caused by the malignant proliferation of bone marrow-derived cells, which is driven by the chromosomal translocation, leading to the fusion of the promyelocytic leukemia gene (*PML*) and the retinoic acid receptor- α gene (*RARA*) (de The et al., 1990; Goto et al., 2011). The oncogenic protein PML-RARA has been found specifically in more than 95% APL patients and is crucial for both the pathogenesis and therapeutics of APL (de The et al., 1990; Zhou et al., 2007; Wang and Chen, 2008). Arsenic trioxide (As_2O_3), an ancient traditional Chinese medicine, has recently been adopted as the first-line treatment for APL patients by the National Comprehensive Cancer Network (Wang and Chen, 2008; Zhu and Huang, 2014), which greatly improved the 5-year overall survival of APL patients (Li et al., 2002; Zhou et al., 2007; Wang and Chen, 2008; Lo-Coco et al., 2013). Mounting biological studies revealed that As_2O_3 cures APL by initiating PML-RARA degradation through specific binding to the PML moiety (Lallemand-Breitenbach et al., 2001; Zhang et al., 2010). Mechanistically, As_2O_3 binds directly to the dicysteine residues C212/213 located in the B2 domain of PML, which induces PML oligomerization, NB formation, SUMOylation, and ultimate degradation of PML (Lallemand-Breitenbach et al., 2008; Jeanne et al., 2010; Zhang et al., 2010).

Unfortunately, although the application of As_2O_3 dramatically promoted the clinical remission of APL patients, some patients still experienced a relapse with As_2O_3 treatment in the clinical practice (Goto et al., 2011; Zhu et al., 2014). Therefore, characterizing the molecular mechanism of As_2O_3 resistance is critical for optimizing the clinical application of As_2O_3 . Recently, it has been reported that 9 arsenic-resistant APL patients harboring PML-RARA mutations, including S214L, A216T, L217F and S220G which locate in the B2 domain of PML-RARA (Zhu et al., 2014). Among them, 8 patients could not be induced a second remission and died (Zhu et al., 2014). These clinical observations provided *in vivo* evidence that mutations in the hot-spot domain (C212-S220) of PML-

RARA might contribute to the unresponsiveness to As_2O_3 treatment, and the underlying molecular mechanism of PML-RARA mutations with As_2O_3 resistance deserve further study.

In this study, we constructed the expression plasmids of PML-RARA-S214L, A216T, L217F and S220G that were clinically observed in arsenic-resistant APL patients (Fig. 1A). APL cell line NB4 cells were transfected with the indicated plasmid and a time-course study was performed with 2 μ mol/L As_2O_3 . As shown in Fig. 1B, As_2O_3 induced a significant degradation of wild-type PML-RARA. Similar results were observed with the mutants L217F and S220G. Conversely, addition of As_2O_3 did not induce the degradation of S214L, even the treatment time was extended to 36 h. While in the case of A216T, the abundance of protein decreased moderately compared with wild-type PML-RARA. Quantification for the protein abundance of distinct PML-RARA mutants with As_2O_3 treatment revealed that mutations occurred in different sites of PML-RARA B2 domain response distinctly to As_2O_3 treatment (Fig. 1C). The unresponsiveness of mutant S214L to As_2O_3 indicates the essential role of Ser214 amino residue in mediating As_2O_3 -induced physiological consequence.

As_2O_3 -induced degradation of PML-RARA is initiated by the direct binding between As_2O_3 and PML-RARA (Jeanne et al., 2010). As different degradation pattern was observed above in the PML-RARA mutants, to detect the binding affinity of As_2O_3 to different PML-RARA mutants, streptavidin agarose affinity assay was performed with *p*-aminophenylarsine oxide (PAPAO) as the probe (Zhang et al., 2010). PAPAO was conjugated to biotin, which was designed as Biotin-As. The result showed that biotin-As bound to PML-RARA, similar binding affinity was observed with PML-RARA-L217F and S220G (Fig. 1D and 1E). And the mutant A216T showed a relatively decreased binding affinity compared with the wild-type PML-RARA. Conversely, S214L mutation exhibited sharply diminished biotin-As binding (Fig. 1D and 1E), similar to the mutant C212A which has been demonstrated to mediate the direct arsenic binding (Jeanne et al., 2010). These results indicate that Ser214, together with the adjacent dicysteine motif C212/213, are




Figure 1. The effects of AS₂O₃ on the degradation of PML-RARA mutants. (A) Schematic model of PML, RARA, and the fusion protein. The functional domains of PML and RARA are indicated with black and gray, respectively. RING refers to the RING finger; B1 and B2, B-box motifs; C-C, coiled-coil; and DBD, DNA-binding domain. Black arrows indicate the break points of the fusion proteins PML-RARA. The CC motif (C212/213) required for AS₂O₃ binding is underlined. The mutation sites Ser214, Ala216, Leu217 and Ser220 were highlighted in bold. (B) NB4 cells transfected with different plasmids were treated with 2 μmol/L AS₂O₃ for the indicated time and AS₂O₃-induced PML-RARA degradation was evaluated by Western blot using anti-Flag antibody. (C) Quantification of the relative protein level of PML-RARA mutants with AS₂O₃ treatment showed in Fig. 1B was calculated. The protein abundance of PML-RARA without AS₂O₃ treatment (0 h) was presented as '1' and the level of PML-RARA with AS₂O₃ treatment for the indicated time was expressed relatively to this. The error bars represent standard error. (D) CHO cells transfected with indicated PML-RARA mutants were treated with 10 μmol/L AS₂O₃ for 2 h. Cells were harvested and lysed in 8 mol/L urea buffer. The supernatant was incubated with streptavidin agarose beads overnight at 4°C. After extensive wash, the streptavidin beads were resuspended in 30 μL SDS-PAGE loading buffer. The binding of arsenic with PML-RARA was analyzed by Western blot. (E) The binding affinity of PML-RARA mutants with arsenic observed in Fig. 1D was quantified. The ratio was described relatively to the binding affinity of wild-type PML-RARA with arsenic.

equally essential in mediating the arsenic-PML binding, consistent with the data that S214L mutant is resistant to AS₂O₃-induced degradation.

Upon the binding of arsenic to PML-RARA, PML-RARA is subjected to hierarchical posttranslational modifications, such as SUMOylation, which is required for the responsiveness of PML-RARA to AS₂O₃ (Lallemand-Breitenbach et al., 2008; Jeanne et al., 2010; Zhang et al., 2010; Goto et al., 2011). Because distinct mutants of PML-RARA exhibited different AS₂O₃ binding affinity and degradation pattern followed by AS₂O₃ treatment, we next detected whether these mutants affect the basal SUMOylation of PML-RARA. As expected, SUMOylated PML bands were observed in cells transfected with wild-type PML-RARA. In contrast, the SUMOylation level of A216T elicited a decreased pattern, while mutation L217F and S220G showed no significant effect on PML-RARA SUMOylation. Notably, no SUMOylation band was observed for S214L (Fig. 2A). To confirm the data of S214L, the SUMOylated PML-RARA bands were detected with AS₂O₃ treatment. Compared with PML-RARA, SUMOylated S214L was not detected with or without AS₂O₃ treatment (Fig. 2B). The varying SUMOylation pattern of PML-RARA mutants is correlated with the degradation pattern of these mutants.


It has recently been demonstrated that AS₂O₃ exposure triggers the nucleoplasmic PML transfers toward the nuclear matrix and nuclear body (NB) (Jeanne et al., 2010; Goto et al., 2011). To detect the transfer of different PML-RARA mutants in response to AS₂O₃, we performed subcellular fraction assay using 293T cells that expressing wild-type PML-RARA or the mutants. As expected, in the presence of AS₂O₃, PML-RARA was observed mostly in the insoluble nuclear body fraction (Fig. 2C, lane 2). Similar results were detected with the mutants of L217F and S220G (Fig. 2C, lanes 8 and 10). Conversely, the transfer of S214L exhibited no response to AS₂O₃ treatment (Fig. 2C, lanes 3 and 4). The relative protein intensity in the soluble or insoluble fraction was quantified and depicted in Fig. 2D. It has been reported that PML-RARA forms multimerization via disulfide-mediated covalent binding in response to AS₂O₃, and subsequently triggers the NB formation (Jeanne et al., 2010). Given that the essential role of Ser214 in AS₂O₃ response, we performed immunofluorescence assay to detect the effect of mutation S214L on the subcellular localization of PML-RARA. HeLa cells were transfected with the indicated PML-RARA expression plasmids and treated with or without AS₂O₃. Wild-type PML-RARA and mutant S220G were detected in the nucleus as a microgranular pattern (Fig. 2E) and the localization was sharply altered to a macrogranular pattern in response to AS₂O₃ (Fig. 2E). Interestingly, S214L was localized diffusely in the nucleus in the presence or

absence of AS₂O₃ (Fig. 2E), suggesting a non-responsiveness of S214L to AS₂O₃ treatment. Previous studies demonstrate that the combination of all-trans retinoic acid (ATRA) works better than AS₂O₃ alone in APL treatment. Therefore, we hypothesized that combining ATRA with AS₂O₃ might induce the degradation of S214L and A216T mutants. As expected, AS₂O₃-ATRA triggered the degradation of S214L and A216T (Fig. 2F). These results suggest that combination of AS₂O₃ with ATRA overcome the arsenic-resistance of PML-RARA mutations.

In this study, we revealed the molecular mechanism of the responsiveness of the recent clinically identified PML-RARA mutations to AS₂O₃ treatment. Among these mutants, S214L strikingly disrupted PML-arsenic binding, NB formation, basal SUMOylation, and degradation. Compared with S214L, mutation A216T showed moderate defect in AS₂O₃ response, while mutants L217F and S220G behave similarly to wild-type PML-RARA (Fig. 2G). These findings suggest distinct mutants of PML-RARA confer varying degree of AS₂O₃ resistance.

Results from GOTO et al. also reported the missense mutations of PML-RARA-A216V and L218P occurred in 2 patients with refractory/relapsed APL (Goto et al., 2011). These two mutations contribute to the aberrant PML-RARA SUMOylation and NB formation (Goto et al., 2011). When our work was underway, Liu et al. also demonstrated that PML-RARA S214L, A216T, L217F and S220G showed varying resistance to AS₂O₃ (Liu et al., 2016). In this study, we detected the direct binding of PML-RARA mutants with AS₂O₃. Mutant S214L exhibited no binding to AS₂O₃, demonstrating that S214, together with C212/213, is equally essential for mediating the effect of AS₂O₃. It is possible that mutations in these amino acid residues lead to a conformational change in PML structure, which disrupts the microenvironment for the direct PML-AS₂O₃ binding and abolishes the following physiological consequences including SUMOylation and degradation of PML-RARA. Further determination of the three dimensional structure of PML-RARA and PML-RARA-S214L are needed to elucidate the detailed molecular mechanism.

AS₂O₃ has been considered as a consolidation treatment for the relapsed APL patients, which has dramatically improved the event-free survival and overall survival in APL patients (Rice and de The, 2014). Mechanism study about the mutations in PML-RARA for AS₂O₃ resistance indicates that higher dosage of AS₂O₃ or the combination of AS₂O₃ with chemotherapy might be helpful for those patients harboring A216T mutation. The unresponsiveness of S214L to AS₂O₃ exposure suggests that it is better to choose an alternative treatment strategy other than AS₂O₃ in relapsed APL patients.



◀ **Figure 2. Ser214 is required for the SUMOylation of PML-RARA.** (A) CHO cells were transfected with His-SUMO2 and PML-RARA or the mutants. At 36 h after transfection, cells were harvested and subjected to *in vivo* SUMOylation analysis. (B) 293T cells were transfected with the indicated plasmids and then cells were incubated with or without with AS₂O₃ for 4 h. (C) 293T cells harboring the expression plasmids of PML-RARA mutants were treated with or without 2 μmol/L AS₂O₃ for 4 h. Cells were harvested and lysed with RIPA buffer, and then the whole cell lysates were separated into soluble and insoluble fractions by centrifugation followed by immunoblotting. (D) The protein intensity of PML-RARA mutants in soluble (S) or insoluble (In) fraction was quantified. The ratio was expressed relatively to the sample without AS₂O₃ treatment. (E) HeLa cells transfected with flag-tagged PML-RARA or mutants S214L and S220G were treated with or without 10 μmol/L AS₂O₃ for 6 h. Anti-Flag antibody was used to detect the NB formation and the nuclear was stained with DAPI. Scale bar, 10 μm. (F) HeLa cells transfected with S214L or A216T expressing plasmids were treated with the drug (2 μmol/L, respectively) for 16 h. The abundance of S214L and A216T was detected by western blot with anti-Flag antibody. (G) On the basis of our and previous data, arsenic directly binds to the Cys212/213 and Ser214 amino acids in the PML B2 domain, leading to the formation of oligomerization and promote SUMOylation of PML, which followed by the ubiquitination and degradation of PML-RARA. Mutation of Ser214 totally disrupts the response of PML-RARA to AS₂O₃. A216T shows moderately defects in the response to AS₂O₃. L217F and S220G, mutations have no effect on the response of PML-RARA to AS₂O₃.

FOOTNOTES

We sincerely thank Prof. Jun Tang for providing the cDNA of PML-RARA, and Dr. Lie Ma for the Biotin and AS₂O₃ used in this study. This work was supported by the National Natural Science Foundation of China (Grant Nos. 31470754 and 31670786), the National High Technology and Development Program of China (2016YFC1302400) and Doctoral Fund of Ministry of Education of China (20130001130003).

D.-M. B and X.-F. Z designed this study. D.-M. B performed the experiments and analyzed the data. D.-M. B and X.-F. Z wrote the paper.

Dong-Mei Bai and Xiao-Feng Zheng declare that they have no conflict of interests. This article does not contain any studies with human or animal subjects performed by the any of the authors.

Dong-Mei Bai^{1,2}, Xiao-Feng Zheng^{1,2}✉

¹ State Key Lab of Protein and Plant Gene Research, Beijing 100871, China

² Department of Biochemistry and Molecular Biology, School of Life Sciences, Peking University, Beijing 100871, China

✉ Correspondence: xiaofengz@pku.edu.cn (X.-F. Zheng)

OPEN ACCESS

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

REFERENCES

- de The H, Chomienne C, Lanotte M, Degos L, Dejean A (1990) The t(15;17) translocation of acute promyelocytic leukaemia fuses the retinoic acid receptor alpha gene to a novel transcribed locus. *Nature* 347:558–561
- Goto E, Tomita A, Hayakawa F, Atsumi A, Kiyoi H, Naoe T (2011) Missense mutations in PML-RARA are critical for the lack of responsiveness to arsenic trioxide treatment. *Blood* 118:1600–1609
- Jeanne M, Lallemand-Breitenbach V, Ferhi O, Koken M, Le Bras M, Duffort S, Peres L, Berthier C, Soilihi H, Raught B et al (2010) PML/RARA oxidation and arsenic binding initiate the antileukemia response of AS₂O₃. *Cancer Cell* 18:88–98
- Lallemand-Breitenbach V, Zhu J, Puvion F, Koken M, Honore N, Doubeikovskiy A, Duprez E, Pandolfi PP, Puvion E, Freemont P et al (2001) Role of promyelocytic leukemia (PML) sumolation in nuclear body formation, 11S proteasome recruitment, and AS₂O₃-induced PML or PML/retinoic acid receptor alpha degradation. *J Exp Med* 193:1361–1371
- Lallemand-Breitenbach V, Jeanne M, Benhenda S, Nasr R, Lei M, Peres L, Zhou J, Zhu J, Raught B, de The H (2008) Arsenic degrades PML or PML-RARalpha through a SUMO-triggered RNF4/ubiquitin-mediated pathway. *Nat Cell Biol* 10:547–555
- Li J, Chen P, Sinogeeva N, Gorospe M, Wersto RP, Chrest FJ, Barnes J, Liu Y (2002) Arsenic trioxide promotes histone H3 phosphoacetylation at the chromatin of CASPASE-10 in acute promyelocytic leukemia cells. *J Biol Chem* 277:49504–49510
- Liu J, Zhu HH, Jiang H, Jiang Q, Huang XJ (2016) Varying responses of PML-RARA with different genetic mutations to arsenic trioxide. *Blood* 127:243–250
- Lo-Coco F, Avvisati G, Vignetti M, Thiede C, Orlando SM, Iacobelli S, Ferrara F, Fazi P, Cicconi L, Di Bona E et al (2013) Retinoic acid and arsenic trioxide for acute promyelocytic leukemia. *N Engl J Med* 369:111–121
- Rice KL, de The H (2014) The acute promyelocytic leukaemia success story: curing leukaemia through targeted therapies. *J Intern Med* 276:61–70
- Wang ZY, Chen Z (2008) Acute promyelocytic leukemia: from highly fatal to highly curable. *Blood* 111:2505–2515
- Zhang XW, Yan XJ, Zhou ZR, Yang FF, Wu ZY, Sun HB, Liang WX, Song AX, Lallemand-Breitenbach V, Jeanne M et al (2010) Arsenic trioxide controls the fate of the PML-RARalpha oncoprotein by directly binding PML. *Science* 328:240–243

Zhou GB, Zhang J, Wang ZY, Chen SJ, Chen Z (2007) Treatment of acute promyelocytic leukaemia with all-trans retinoic acid and arsenic trioxide: a paradigm of synergistic molecular targeting therapy. *Philos Trans R Soc Lond B Biol Sci* 362:959–971

Zhu HH, Huang XJ (2014) Oral arsenic and retinoic acid for non-high-risk acute promyelocytic leukemia. *N Engl J Med* 371:2239–2241

Zhu HH, Qin YZ, Huang XJ (2014) Resistance to arsenic therapy in acute promyelocytic leukemia. *N Engl J Med* 370:1864–1866

Electronic supplementary material The online version of this article (doi:[10.1007/s13238-016-0356-4](https://doi.org/10.1007/s13238-016-0356-4)) contains supplementary material, which is available to authorized users.